

# THE HVZ THEOREM FOR A PSEUDO-RELATIVISTIC OPERATOR

D. H. JAKUBASSA-AMUNDSEN

ABSTRACT. The localization of the essential spectrum of a relativistic two-electron ion is provided. The analysis is performed with the help of the pseudo-relativistic Brown-Ravenhall operator which is the restriction of the Coulomb-Dirac operator to the electrons' positive spectral subspace.

## 1. INTRODUCTION

We consider two interacting electrons in a central Coulomb field, generated by a point nucleus of charge number  $Z$  and infinite mass. In contrast to the case of a single electron in the central field, the additional electron-electron interaction potential does not vanish at infinity if simultaneously, the distance between the two electrons is kept fixed. Therefore, the determination of the essential spectrum with the help of relative compactness arguments is not possible. In order to cope with this difficulty, two-cluster decompositions of the involved particles (including the nucleus) are made and a subordinate partition of unity is defined. This method is described in [4, §3] and [21] for the Schrödinger case and is applied by Lewis, Siedentop and Vugalter [13] to the scalar pseudo-relativistic Hamiltonian which is obtained from the Schrödinger operator by replacing the single-particle kinetic energy operator  $p^2/2m$  with  $\sqrt{p^2 + m^2} - m$ ,  $\mathbf{p}$  and  $m$  being the momentum and mass of the electron, respectively, and which has been analyzed in [25] and [9]. Historically, the location of the essential spectrum of multiparticle Schrödinger operators, called HVZ theorem, was provided by Hunziker (using diagrammatic techniques [10]), van Winter [26] and Zhislin [28] (see e.g. [18, p.120,343]). Alternative methods for the determination of the essential spectrum of generalized Schrödinger operators involve  $C^*$ -algebra techniques (see e.g. [7] and references therein).

The Brown-Ravenhall operator to be discussed below was introduced [2] as projection of the Coulomb-Dirac operator onto the positive spectral subspace of the free electrons, and was analyzed in a series of papers (e.g. [6, 23, 24, 1]). It also emerges as the first-order term in unitary transformation schemes applied to the Coulomb-Dirac operator [5, 11]. Such transformations allow for the decoupling of the electron and positron spectral subspaces to arbitrary order  $n$  in the potential

---

*Date:* November 23, 2005.

strength. The convergence of the resulting series of operators was shown only recently [19].

Let (in relativistic units,  $\hbar = c = 1$ )

$$H = \sum_{k=1}^2 (D_0^{(k)} + V^{(k)}) \quad (1.1)$$

be the Dirac operator for two noninteracting electrons, where  $D_0^{(k)} := \boldsymbol{\alpha}^{(k)} \mathbf{p}_k + \beta^{(k)} m$  (with  $\mathbf{p} := -i\nabla_{\mathbf{x}}$ ) is the free Dirac operator [22], and  $V^{(k)} := -\frac{\gamma}{x_k}$  (with  $\gamma = Ze^2$ ,  $e^2 \approx 1/137.04$  the fine-structure constant and  $x := |\mathbf{x}|$ ) is the central Coulomb potential for electron  $k$ .  $H$  acts in the Hilbert space  $\mathcal{A}(L_2(\mathbb{R}^3) \otimes \mathbb{C}^4)^2$  (we shall use this notation as shorthand for  $\mathcal{A}(\bigotimes_{k=1}^2 (L_2(\mathbb{R}^3) \otimes \mathbb{C}^4)^{(k)})$  where  $\mathcal{A}$  denotes antisymmetrization of the two-electron function, and the form domain is the subspace  $\mathcal{A}(H_{1/2}(\mathbb{R}^3) \otimes \mathbb{C}^4)^2$ ).

Let  $V^{(12)} := \frac{e^2}{|\mathbf{x}_1 - \mathbf{x}_2|}$  be the electron-electron interaction. Then the (two-particle) Brown-Ravenhall operator is defined by

$$\begin{aligned} H^{BR} &= \Lambda_{+,2} (H + V^{(12)}) \Lambda_{+,2} \\ &= \frac{1 + \beta^{(1)}}{2} \frac{1 + \beta^{(2)}}{2} U_0^{(1)} U_0^{(2)} (H + V^{(12)}) (U_0^{(1)} U_0^{(2)})^{-1} \frac{1 + \beta^{(1)}}{2} \frac{1 + \beta^{(2)}}{2} \end{aligned} \quad (1.2)$$

where  $\Lambda_{+,2} = \Lambda_+^{(1)} \Lambda_+^{(2)}$  (as short-hand for  $\Lambda_+^{(1)} \otimes \Lambda_+^{(2)}$ ) is the (tensor) product of the single-particle projectors  $\Lambda_+^{(k)}$  onto the positive spectral subspace of  $D_0^{(k)}$ , and the second equality results from the representation of a single-particle function  $\varphi_+^{(k)}$  in this subspace in terms of the Foldy-Wouthuysen transformation  $U_0^{(k)}$ , viz.  $\varphi_+^{(k)} = U_0^{(k)-1} \begin{pmatrix} u_+ \\ 0 \end{pmatrix} = U_0^{(k)-1} \frac{1 + \beta^{(k)}}{2} \begin{pmatrix} u_+ \\ u_- \end{pmatrix}$ , with  $\beta = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}$ ,  $I \in \mathbb{C}^{2,2}$  the unit matrix,  $u_+, u_- \in H_{1/2}(\mathbb{R}^3) \otimes \mathbb{C}^2$ , and

$$U_0^{(k)} := A(p_k) + \beta^{(k)} \boldsymbol{\alpha}^{(k)} \mathbf{p}_k g(p_k), \quad (1.3)$$

$$A(p) := \left( \frac{E_p + m}{2E_p} \right)^{1/2}, \quad g(p) := \frac{1}{\sqrt{2E_p(E_p + m)}}$$

where  $E_p = |D_0| = \sqrt{p^2 + m^2}$ . The inverse  $U_0^{(k)-1} = A(p_k) + \boldsymbol{\alpha}^{(k)} \mathbf{p}_k g(p_k) \beta^{(k)}$ . Since  $H^{BR}$  is sandwiched between the projectors  $\frac{1 + \beta^{(k)}}{2}$  it has a block-diagonal form with one nonvanishing entry,  $h^{BR}$ , defined by means of [6]

$$(\phi_+, H^{BR} \phi_+) = (u, h^{BR} u) \quad (1.4)$$

with  $\phi_+ \in \Lambda_{+,2}(\mathcal{A}(H_{1/2}(\mathbb{R}^3) \otimes \mathbb{C}^4)^2)$  and  $u \in \mathcal{A}(H_{1/2}(\mathbb{R}^3) \otimes \mathbb{C}^2)^2$ , thus reducing the single-particle spinor degrees of freedom from 4 to 2. One obtains (see e.g. [6])

$$h^{BR} = \sum_{k=1}^2 (T^{(k)} + b_{1m}^{(k)}) + v^{(12)} \quad (1.5)$$

$$T^{(k)} := E_{p_k}, \quad b_{1m}^{(k)} := -\gamma \left( A(p_k) \frac{1}{x_k} A(p_k) + \boldsymbol{\sigma}^{(k)} \mathbf{p}_k g(p_k) \frac{1}{x_k} g(p_k) \boldsymbol{\sigma}^{(k)} \mathbf{p}_k \right)$$

where  $\boldsymbol{\sigma}^{(k)}$  is the vector of Pauli matrices, while  $v^{(12)}$  results from the electron-electron interaction term and is specified later (section 5).

We note that  $h^{BR}$  is a well-defined (in the form sense), positive operator for potential strengths  $\gamma \leq \gamma_{BR} = \frac{2}{\pi/2+2/\pi} \approx 0.906$  which relies on the estimates  $(u, \sum_{k=1}^2 (T^{(k)} + b_{1m}^{(k)}) u) \geq 2m(1-\gamma)(u, u)$  [24] and  $V^{(12)} \geq 0$ . In particular from the respective property of the single-particle operator [6] and using [3, 11] that  $(\phi_+, V^{(12)} \phi_+) \leq \frac{e^2}{2\gamma_{BR}} (\phi_+, \sum_{k=1}^2 T^{(k)} \phi_+)$ , the total potential  $V := b_{1m}^{(1)} + b_{1m}^{(2)} + v^{(12)}$  is  $(T^{(1)} + T^{(2)})$ -form bounded with form bound less than one for  $\gamma < \gamma_{BR}$ . Thus  $h^{BR}$  is a self-adjoint operator by means of the Friedrichs extension of the restriction of  $h^{BR}$  to  $\mathcal{A}(C_0^\infty(\mathbb{R}^3) \otimes \mathbb{C}^2)^2$ .

## 2. THE HVZ THEOREM AND THE STRATEGY OF PROOF

We introduce the three two-cluster decompositions of our operator,

$$h^{BR} = T + a_j + r_j, \quad j = 0, 1, 2, \quad (2.1)$$

where  $T := T^{(1)} + T^{(2)}$  and  $a_j$  collects all interactions not involving particle  $j$  ( $j = 1, 2$  denotes the two electrons and  $j = 0$  refers to the nucleus which is fixed at the origin). The remainder  $r_j$  is supposed to vanish when particle  $j$  is moved to infinity (respectively both electrons are moved to infinity in the case  $j = 0$ ). Define for  $j = 0, 1, 2$ ,

$$\Sigma_0 := \min_j \inf \sigma(T + a_j) = \min \{ \inf \sigma(T + b_{1m}^{(1)}), \inf \sigma(T + v^{(12)}) \} \quad (2.2)$$

(note that the two electrons move in the same potential, such that  $b_{1m}^{(1)}(\mathbf{x}_1) = b_{1m}^{(2)}(\mathbf{x}_2)$ ). Then we have

**Theorem 1** (HVZ theorem).

*Let  $h^{BR} = T + b_{1m}^{(1)} + b_{1m}^{(2)} + v^{(12)}$  be the two-electron Brown-Ravenhall operator with potential strength  $\gamma < \gamma_{BR}$ , and let (2.1) be its two-cluster decompositions. Then the essential spectrum of  $h^{BR}$  is given by*

$$\sigma_{ess}(h^{BR}) = [\Sigma_0, \infty). \quad (2.3)$$

Physically this means that the bottom of the essential spectrum is given by the ground state of the one-electron ion, increased by the rest mass of the second electron.

The strategy of proof is based on Simon [20] (see also [4]) as well as on Lieb and Yau [14] and Lewis et al [13]. We start by characterizing the essential spectrum by means of a Weyl sequence, located outside a ball  $B_n(0) \subset \mathbb{R}^6$  of radius  $n$  centered at the origin.

**Lemma 1.** *Let  $h^{BR} = T + V$ , let  $V$  be relatively form bounded with respect to  $T$ . Then  $\lambda \in \sigma_{ess}(h^{BR})$  iff there exists a sequence of functions  $\varphi_n \in \mathcal{A}(C_0^\infty(\mathbb{R}^6 \setminus B_n(0)) \otimes \mathbb{C}^4)$  with  $\|\varphi_n\| = 1$  such that*

$$\|(h^{BR} - \lambda) \varphi_n\| \longrightarrow 0 \quad \text{as } n \rightarrow \infty. \quad (2.4)$$

Recall that  $\varphi_n$  is a two-particle function, each particle being described by a two-spinor. For Schrödinger operators, Lemma 1 is proven in [4, Thm 3.11]. We note that Lemma 1 holds also for the single-particle operators,  $h := T^{(1)}$  or  $h := T^{(1)} + b_{1m}^{(1)}$ , with a proof closely following the one given in section 7.

**Lemma 2** (Persson's theorem).

*Let  $h^{BR} = T + V$ , let  $V$  be relatively form bounded with respect to  $T$ , and let  $\varphi \in \mathcal{A}(C_0^\infty(\mathbb{R}^6 \setminus B_R(0)) \otimes \mathbb{C}^4)$ . Then*

$$\inf \sigma_{ess}(h^{BR}) = \lim_{R \rightarrow \infty} \inf_{\|\varphi\|=1} (\varphi, h^{BR} \varphi). \quad (2.5)$$

The proof given in [4, Thm 3.12] relies on Lemma 1 and on the min-max principle [18, XIII.1], [12, p.60]. It also holds in our case.

Let us now introduce the Ruelle-Simon partition of unity  $(\phi_j)_{j=0,1,2} \in C^\infty(\mathbb{R}^6)$  which is subordinate to the cluster decomposition (2.1), see e.g. [4, p.33], [21]. It is defined on the unit sphere and has the following properties,

$$\sum_{j=0}^2 \phi_j^2 = 1, \quad \phi_j(\lambda \mathbf{x}) = \phi_j(\mathbf{x}) \quad \text{for } x = 1 \text{ and } \lambda \geq 1,$$

$\text{supp } \phi_j \cap \mathbb{R}^6 \setminus B_1(0) \subseteq \{\mathbf{x} \in \mathbb{R}^6 \setminus B_1(0) : |\mathbf{x}_1 - \mathbf{x}_2| \geq Cx \text{ and } x_j \geq Cx\}, \quad j = 1, 2,$

$$\text{supp } \phi_0 \cap \mathbb{R}^6 \setminus B_1(0) \subseteq \{\mathbf{x} \in \mathbb{R}^6 \setminus B_1(0) : x_k \geq Cx \quad \forall k \in \{1, 2\}\}, \quad (2.6)$$

where  $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2)$ ,  $x = |\mathbf{x}|$  and  $C$  is a positive constant.

According to Lemma 2, we can always assume  $\varphi \in \mathcal{A}(C_0^\infty(\mathbb{R}^6 \setminus B_R(0)) \otimes \mathbb{C}^4)$  in the following. For later use we introduce a smooth auxiliary function  $\chi \in C^\infty(\mathbb{R}^6)$ ,  $\text{ran } \chi = [0, 1]$ , which is unity on the support of  $\varphi$ . Then,  $\chi\varphi = \varphi$ . Having in mind (2.2) we aim at a localization formula for our operator. We write

$$\begin{aligned} (\varphi, h^{BR} \varphi) &= \left( \sum_{j=0}^2 \phi_j \chi \phi_j \varphi, h^{BR} \varphi \right) \\ &= \sum_{j=0}^2 (\phi_j \varphi, h^{BR} \chi \phi_j \varphi) - \sum_{j=0}^2 (\phi_j \varphi, [h^{BR}, \chi \phi_j] \varphi) \end{aligned} \quad (2.7)$$

where  $[B, A] = BA - AB$  denotes the commutator. One can show that not only the contribution of  $r_j$  to the first term of (2.7) vanishes uniformly as  $R \rightarrow \infty$ , but also the second term containing the commutator. More precisely, one has

**Lemma 3.** *Let  $h^{BR} = T + a_j + r_j$ ,  $(\phi_j)_{j=0,1,2}$  the Ruelle-Simon partition of unity and  $\varphi \in \mathcal{A}(C_0^\infty(\mathbb{R}^6 \setminus B_R(0)) \otimes \mathbb{C}^4)$  with  $R > 1$ . Then*

$$|(\phi_j \varphi, r_j \phi_j \varphi)| \leq \frac{c}{R} \|\varphi\|^2, \quad j = 0, 1, 2, \quad (2.8)$$

where  $c$  is some constant.

We call an operator  $\mathcal{O}$   $\frac{1}{R}$ -bounded if  $\mathcal{O}$  is bounded by  $\frac{c}{R}$ . Thus Lemma 3 states that  $\phi_j r_j \phi_j$  is  $\frac{1}{R}$ -bounded.

**Lemma 4.** *Assume  $h^{BR}$ ,  $\phi_j$  and  $\varphi$  as in Lemma 3,  $R > 2$ . Then*

$$\begin{aligned} (a) \quad & \left| \sum_{j=0}^2 (\phi_j \varphi, [T, \phi_j] \varphi) \right| \leq \frac{c}{R^2} \|\varphi\|^2 \\ (b) \quad & |(\phi_j \varphi, [b_{1m}^{(k)}, \phi_j] \varphi)| \leq \frac{c}{R} \|\varphi\|^2 \\ (c) \quad & |(\phi_j \varphi, [v^{(12)}, \phi_j] \varphi)| \leq \frac{c}{R} \|\varphi\|^2 \end{aligned} \quad (2.9)$$

where  $c$  is a generic constant.

The proof of (a) in Lemma 4 is provided in [13].

With Lemmata 3 and 4, (2.7) turns into the localization formula

$$(\varphi, h^{BR} \varphi) = \sum_{j=0}^2 (\phi_j \varphi, (T + a_j) \phi_j \varphi) + O\left(\frac{1}{R}\right) \|\varphi\|^2 \quad (2.10)$$

for  $R > 2$ . Using Persson's theorem (Lemma 2), we obtain

$$\inf \sigma_{ess}(h^{BR}) = \lim_{R \rightarrow \infty} \inf_{\|\varphi\|=1} \sum_{j=0}^2 (\phi_j \varphi, (T + a_j) \phi_j \varphi). \quad (2.11)$$

Recalling the definition (2.2) of  $\Sigma_0$  in terms of the smallest infimum of  $\sigma(T + a_j)$ , we can estimate

$$\inf_{\|\varphi\|=1} \sum_{j=0}^2 (\phi_j \varphi, (T + a_j) \phi_j \varphi) \geq \sum_{j=0}^2 \Sigma_0 (\phi_j \varphi, \phi_j \varphi) = \Sigma_0 \quad (2.12)$$

since  $\sum_{j=0}^2 \phi_j^2 = 1 = \|\varphi\|^2$ . This proves the inclusion  $\sigma_{ess}(h^{BR}) \subset [\Sigma_0, \infty)$ .

For the remaining inclusion,  $[\Sigma_0, \infty) \subset \sigma_{ess}(h^{BR})$ , in the literature called the 'easy part' of the proof of the HVZ theorem, we use the strategy of Weyl sequences [4, 21].

Let  $\lambda \in [\Sigma_0, \infty)$ . Consider the case that  $\Sigma_0 = \inf \sigma(T + a_j)$  for  $j = 1$ , and assume that  $\lambda \in \sigma(T + a_1)$ . (This assumption is proven in (2.16) where it is shown that  $\sigma(T + a_1) = \sigma_{ess}(T + a_1)$ .) Since  $T + a_1 = T^{(1)} + (T^{(2)} + b_{1m}^{(2)})$  describes two independent particles, we can decompose  $\lambda = \lambda_1 + \lambda_2$  with  $\lambda_1 \in \sigma_{ess}(T^{(1)})$  and  $\lambda_2 \in \sigma(T^{(2)} + b_{1m}^{(2)})$ .

Let  $(\varphi_n^{(1)})_{n \in \mathbb{N}}$  be a Weyl sequence corresponding to  $\lambda_1$ , i.e.  $\varphi_n^{(1)}$  is characterized by

$$\|\varphi_n^{(1)}\| = 1, \quad \varphi_n^{(1)} \xrightarrow{w} 0, \quad \|(T^{(1)} - \lambda_1) \varphi_n^{(1)}\| \rightarrow 0 \text{ as } n \rightarrow \infty. \quad (2.13)$$

From Lemma 1 we can require  $\varphi_n^{(1)} \in C_0^\infty(\mathbb{R}^3 \setminus B_n(0)) \otimes \mathbb{C}^2$ . Let  $(\phi_n^{(2)})_{n \in \mathbb{N}} \in C_0^\infty(\mathbb{R}^3) \otimes \mathbb{C}^2$  be a defining sequence for  $\lambda_2$  according to [27, Thm 7.22] with

$\|\phi_n^{(2)}\| = 1$ . Since  $\|(T^{(2)} + b_{1m}^{(2)} - \lambda_2)\phi_n^{(2)}\| \rightarrow 0$  as  $n \rightarrow \infty$ , for any given  $\epsilon > 0$  there exists  $N \in \mathbb{N}$  such that

$$\|(T^{(2)} + b_{1m}^{(2)} - \lambda_2)\phi_N^{(2)}\| < \epsilon. \quad (2.14)$$

We define the sequence  $\psi_n := \varphi_n^{(1)} \cdot \phi_N^{(2)}$  for  $n \in \mathbb{N}$  and the antisymmetric sequence  $\mathcal{A}\psi_n := \frac{1}{\sqrt{2}}(\varphi_n^{(1)}\phi_N^{(2)} - \varphi_n^{(2)}\phi_N^{(1)})$  and claim that a subsequence of  $\mathcal{A}\psi_n$  is a Weyl sequence for  $\lambda \in \sigma_{ess}(h^{BR})$ .

(i) The weak convergence,  $\mathcal{A}\psi_n \xrightarrow{w} 0$ , follows from

$$\begin{aligned} \sqrt{2} |(\mathcal{A}\psi_n, f^{(1)}g^{(2)})| &\leq |(\varphi_n^{(1)}, f^{(1)})| \cdot |(\phi_N^{(2)}, g^{(2)})| + |(\phi_N^{(1)}, f^{(1)})| \cdot |(\varphi_n^{(2)}, g^{(2)})| \\ &\leq |(\varphi_n^{(1)}, f^{(1)})| \|\phi_N^{(2)}\| \|g^{(2)}\| + |(\varphi_n^{(2)}, g^{(2)})| \|\phi_N^{(1)}\| \|f^{(1)}\| \rightarrow 0 \end{aligned} \quad (2.15)$$

for all  $f^{(1)}, g^{(2)} \in L_2(\mathbb{R}^3) \otimes \mathbb{C}^2$ , since by (2.13)  $\varphi_n^{(1)} \xrightarrow{w} 0$ .

(ii)  $(\psi_n)_{n \in \mathbb{N}}$  obeys the Weyl criterion for  $\lambda \in \sigma_{ess}(T + a_1)$  since

$$\|(T^{(1)} - \lambda_1 + T^{(2)} + b_{1m}^{(2)} - \lambda_2)\psi_n\| \quad (2.16)$$

$$\leq \|(T^{(1)} - \lambda_1)\varphi_n^{(1)}\| \|\phi_N^{(2)}\| + \|(T^{(2)} + b_{1m}^{(2)} - \lambda_2)\phi_N^{(2)}\| \|\varphi_n^{(1)}\| < 2\epsilon$$

for arbitrary  $\epsilon$  and  $n$  sufficiently large because of (2.13) and (2.14).

(iii) Using that  $h^{BR}$  is symmetric upon particle exchange, we have

$$\|(h^{BR} - \lambda)\mathcal{A}\psi_n\| \leq 2\|(h^{BR} - \lambda)\frac{1}{\sqrt{2}}\varphi_n^{(1)}\phi_N^{(2)}\|$$

$$\leq \sqrt{2}\|(T + a_1 - \lambda)\psi_n\| + \sqrt{2}\|r_1\psi_n\| < 2\sqrt{2}\epsilon + \sqrt{2}\|b_{1m}^{(1)}\psi_n\| + \sqrt{2}\|v^{(12)}\psi_n\| \quad (2.17)$$

where (2.16) was used. One can show that the two remaining terms also tend to zero as  $n \rightarrow \infty$ . More precisely, one has

**Lemma 5.** *Let  $\varphi \in C_0^\infty(\mathbb{R}^3 \setminus B_R(0)) \otimes \mathbb{C}^2$  and  $R > 1$ . Then for some constant  $c$ ,*

$$\|b_{1m}^{(1)}\varphi\| \leq \frac{c}{R}\|\varphi\|. \quad (2.18)$$

**Lemma 6.** *Let  $\psi_n$  as defined above. Then for all  $\varphi \in (C_0^\infty(\mathbb{R}^3) \otimes \mathbb{C}^2)^2$ ,*

$$|(\varphi, v^{(12)}\psi_n)| \leq \frac{c}{n}\|\varphi\|\|\psi_n\| \quad (2.19)$$

with some constant  $c$ .

With Lemma 5, one has  $\|b_{1m}^{(1)}\psi_n\| = \|b_{1m}^{(1)}\varphi_n^{(1)}\| \|\phi_N^{(2)}\| \leq \frac{c}{n}\|\varphi_n^{(1)}\| \|\phi_N^{(2)}\|$ . Moreover, one has the equivalence for an essentially self-adjoint operator  $A$  and  $\psi \in \mathcal{D}(A)$  [16, p.260]

$$(i) \quad \|A\psi\| \leq \frac{c}{n}\|\psi\| \quad (2.20)$$

$$(ii) \quad |(\varphi, A\psi)| \leq \frac{c}{n}\|\varphi\|\|\psi\| \quad \forall \varphi \text{ in the core of } \mathcal{D}(A).$$

Choosing  $A := v^{(12)}$ ,  $\psi := \psi_n$  and  $\varphi \in (C_0^\infty(\mathbb{R}^3) \otimes \mathbb{C}^2)^2$  and using Lemma 6, this proves that the r.h.s. of (2.17) is smaller than  $4\epsilon$  for sufficiently large  $n$ .

(iv) Concerning the normalizability of  $\mathcal{A}\psi_n$ , we have

$$2 \|\mathcal{A}\psi_n\|^2 = \|\varphi_n^{(1)}\| \|\phi_N^{(2)}\| + 2 \operatorname{Re} (\varphi_n^{(1)} \phi_N^{(2)}, \varphi_n^{(2)} \phi_N^{(1)}) + \|\varphi_n^{(2)}\| \|\phi_N^{(1)}\|. \quad (2.21)$$

Since  $\phi_N^{(1)} \in C_0^\infty(\mathbb{R}^3) \otimes \mathbb{C}^2$  there exists an  $R_0 > 0$  such that  $x_1 < R_0$  on  $\operatorname{supp} \phi_N^{(1)}$ , and in addition,  $x_1 > n$  on  $\operatorname{supp} \varphi_n^{(1)}$ . Hence we have

$$(\varphi_n^{(1)} \phi_N^{(2)}, \varphi_n^{(2)} \phi_N^{(1)}) = \left| \int_{\mathbb{R}^3} d\mathbf{x}_1 \overline{\varphi_n^{(1)}} \phi_N^{(1)} \right|^2 = 0 \quad \text{if } n > R_0. \quad (2.22)$$

Thus  $\|\mathcal{A}\psi_n\| = 1$  for sufficiently large  $n$ . Therefore a subsequence of  $\mathcal{A}\psi_n$  is a Weyl sequence for  $\lambda$ , resulting in  $\lambda \in \sigma_{\text{ess}}(h^{BR})$ .

Due to the symmetry upon particle exchange, this proves the case  $j = 2$  as well. Consideration of the case  $j = 0$  can be omitted, since  $V^{(12)} \geq 0$  and hence  $v^{(12)} \geq 0$  (whereas  $V^{(1)} \leq 0$  and so  $b_{1m}^{(1)} \leq 0$ ). Thus  $\inf \sigma(T + v^{(12)}) \geq \inf \sigma(T + b_{1m}^{(1)})$  such that one has  $\Sigma_0 = \inf \sigma(T + b_{1m}^{(1)})$ .

### 3. INGREDIENTS FOR THE PROOFS OF THE LEMMATA

The main difference in the proofs of Lemmata 1, 3, 5 and 6 as contrasted to the proof of Lemma 4 lies in the fact that the momentum representation is used for the former, whereas the proof of Lemma 4 is carried out in coordinate space.

An important estimate which holds in either space is the Lieb and Yau formula (which is related to the Schur test for the boundedness of integral operators and is easily derived from the Schwarz inequality), generalized to the two-particle case [14, 11].

**Lemma 7** (Generalized Lieb and Yau formula).

Let  $A$  be an essentially self-adjoint integral operator and  $k_A(\boldsymbol{\xi}, \boldsymbol{\xi}')$  its kernel,  $\boldsymbol{\xi} \in \mathbb{R}^{3l}$  with  $l \in \{1, 2\}$  denoting the number of particles. Then for  $\psi, \varphi \in \mathcal{D}(A)$ ,

$$\begin{aligned} |(\psi, A\varphi)| &= \left| \int d\boldsymbol{\xi} \overline{\psi(\boldsymbol{\xi})} \int d\boldsymbol{\xi}' k_A(\boldsymbol{\xi}, \boldsymbol{\xi}') \varphi(\boldsymbol{\xi}') \right| \\ &\leq \left( \int d\boldsymbol{\xi} |\psi(\boldsymbol{\xi})|^2 I(\boldsymbol{\xi}) \right)^{\frac{1}{2}} \cdot \left( \int d\boldsymbol{\xi}' |\varphi(\boldsymbol{\xi}')|^2 J(\boldsymbol{\xi}') \right)^{\frac{1}{2}} \end{aligned} \quad (3.1)$$

with

$$\begin{aligned} I(\boldsymbol{\xi}) &:= \int d\boldsymbol{\xi}' |k_A(\boldsymbol{\xi}, \boldsymbol{\xi}')| \frac{f(\boldsymbol{\xi})}{f(\boldsymbol{\xi}')} \\ J(\boldsymbol{\xi}') &:= \int d\boldsymbol{\xi} |k_A(\boldsymbol{\xi}, \boldsymbol{\xi}')| \frac{f(\boldsymbol{\xi}')}{f(\boldsymbol{\xi})}, \end{aligned} \quad (3.2)$$

where  $f(\boldsymbol{\xi})$  is a nonnegative convergence generating function and all integrals run over  $\mathbb{R}^{3l}$ .

If  $|k_A|$  is symmetric in  $\boldsymbol{\xi}, \boldsymbol{\xi}'$  then  $J(\boldsymbol{\xi}') = I(\boldsymbol{\xi})$ . Provided we can show that  $I$  and  $J$  are  $\frac{1}{R}$ -bounded functions (for all values of  $\boldsymbol{\xi}$  and  $\boldsymbol{\xi}'$ ), then the uniform  $\frac{1}{R}$ -boundedness of  $A$  follows from (3.1) and from the equivalence (2.20),

$$\begin{aligned} |(\psi, A \varphi)| &\leq \left( \int d\boldsymbol{\xi} |\psi(\boldsymbol{\xi})|^2 \frac{c}{R} \right)^{\frac{1}{2}} \left( \int d\boldsymbol{\xi}' |\varphi(\boldsymbol{\xi}')|^2 \frac{c}{R} \right)^{\frac{1}{2}} \\ &= \frac{c}{R} \|\psi\| \|\varphi\|, \end{aligned} \quad (3.3)$$

with  $c$  a suitable constant.

Let us now consider the properties of the smooth auxiliary function  $\chi \in C^\infty(\mathbb{R}^{3l})$  where  $l = 2$  denotes the two-particle and  $l = 1$  the one-particle case. For  $\varphi \in C_0^\infty(\mathbb{R}^{3l} \setminus B_R(0)) \otimes \mathbb{C}^{2l}$  we define for  $\mathbf{x} \in \mathbb{R}^{3l}$

$$\chi\left(\frac{\mathbf{x}}{R}\right) := \begin{cases} 0, & x < R/2 \\ 1, & x \geq R \end{cases} \quad (3.4)$$

such that  $\chi = 1$  on  $\text{supp } \varphi$ . Moreover, define

$$\chi_0\left(\frac{\mathbf{x}}{R}\right) := 1 - \chi\left(\frac{\mathbf{x}}{R}\right) \quad (3.5)$$

with  $\chi_0 \in \mathcal{S}(\mathbb{R}^{3l})$ ,  $\mathcal{S}$  being the Schwartz space.

In our proofs we shall introduce commutators with  $\chi$ , such that  $\chi$  can be replaced by  $\chi_0$ , viz.  $[B, \chi] = -[B, \chi_0]$ , and for the operator  $B$  we shall choose a multiplication operator in momentum space. Then one can readily work in Fourier space since the Fourier transform of the Schwartz function  $\chi_0$  is again a Schwartz function, making the resulting integrals converge. Marking the Fourier transform with a hat, we have for  $\mathbf{p} \in \mathbb{R}^{3l}$ ,

$$\begin{aligned} \left( \widehat{\chi_0\left(\frac{\cdot}{R}\right)} \right) (\mathbf{p}) &= \frac{1}{(2\pi)^{3l/2}} \int_{\mathbb{R}^{3l}} d\mathbf{x} e^{-i\mathbf{p}\mathbf{x}} \chi_0\left(\frac{\mathbf{x}}{R}\right) \\ &= \frac{1}{(2\pi)^{3l/2}} R^{3l} \int_{\mathbb{R}^{3l}} d\mathbf{z} e^{-i\mathbf{p}R\mathbf{z}} \chi_0(\mathbf{z}) = R^{3l} \hat{\chi}_0(\mathbf{p}R). \end{aligned} \quad (3.6)$$

In Lemma 4 the commutator with the partition of unity,  $\phi_j$ , is needed. As neither  $\phi_j$  nor  $1 - \phi_j$  is a Schwartz (or even an  $L_2$ ) function, its Fourier transform is not well defined. Therefore we work in coordinate space instead. The strategy we apply is to construct commutators  $[B, \phi_j]$  where  $B$  is again a multiplication operator in momentum space. Then its kernel  $k_B$  factorizes,  $k_B(\mathbf{p}, \mathbf{p}') = B(\mathbf{p}) \delta(\mathbf{p} - \mathbf{p}')$ . Our aim is to estimate this kernel in coordinate space and then apply the Lieb and Yau formula.

Consider the one-particle case,  $\mathbf{p} \in \mathbb{R}^3$ , and let  $B(\mathbf{p}) := (\boldsymbol{\sigma}\mathbf{p})^l g(p)$  where  $g$  is spherically symmetric and  $l \in \{0, 1\}$ . The Fourier transformed kernel is defined by

$$\check{k}_B(\mathbf{x}_1, \mathbf{x}'_1) = \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} d\mathbf{p} e^{i\mathbf{p}\mathbf{x}_1} \int_{\mathbb{R}^3} d\mathbf{p}' e^{-i\mathbf{p}'\mathbf{x}'_1} (\boldsymbol{\sigma}\mathbf{p})^l g(p) \delta(\mathbf{p} - \mathbf{p}')$$



$$\begin{aligned}
&= (-i\boldsymbol{\sigma}\nabla_{\mathbf{x}_1})^l \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} d\mathbf{p} e^{ip(\mathbf{x}_1 - \mathbf{x}'_1)} g(p) \\
&= (-i\boldsymbol{\sigma}\nabla_{\tilde{\mathbf{x}}})^l \frac{4\pi}{(2\pi)^3} \frac{1}{\tilde{x}} \int_0^\infty p dp \sin p\tilde{x} g(p)
\end{aligned} \tag{3.7}$$

where  $\tilde{\mathbf{x}} := \mathbf{x}_1 - \mathbf{x}'_1$  is introduced. This integral can be estimated with the help of complex and asymptotic analysis [15].

In the following we prove successively Lemma 5 (section 4), Lemma 6 (section 5), Lemma 3 (section 6), Lemma 1 (section 7) and Lemma 4 (section 8).

#### 4. PROOF OF LEMMA 5

According to the equivalence (2.20) we prove  $|(\phi, b_{1m}^{(1)}\varphi)| \leq \frac{c}{R} \|\phi\| \|\varphi\|$  for all  $\phi \in C_0^\infty(\mathbb{R}^3) \otimes \mathbb{C}^2$ .

The operator  $b_{1m}^{(1)}$ , defined in (1.5) with (1.3), is a sum of terms each of which has the structure  $B(\mathbf{p})\frac{1}{x}B(\mathbf{p})$  where the indices on  $\mathbf{p}_1$  and  $\mathbf{x}_1$  are suppressed and  $B$  is an analytic (for  $m \neq 0$ ), bounded multiplication operator in momentum space. Hence one can apply the mean value theorem to find

$$B(\mathbf{p}) = B(\mathbf{p}') + (\mathbf{p} - \mathbf{p}') \nabla_{\mathbf{p}} B(\boldsymbol{\xi}_0), \quad \boldsymbol{\xi}_0 := \lambda\mathbf{p} + (1 - \lambda)\mathbf{p}' \tag{4.1}$$

for a suitable  $\lambda \in [0, 1]$ . Since  $B$  is bounded for all  $\mathbf{p}$ , its derivative can be estimated by

$$|\nabla_{\mathbf{p}} B(\mathbf{p})| \leq \frac{c_0}{1+p} \tag{4.2}$$

with some constant  $c_0$ . We introduce the auxiliary function  $\chi$  from (3.4) (with  $l = 1$ ) and estimate (by means of the triangle inequality) each term of  $b_{1m}^{(1)}$  separately,

$$|(\phi, B \frac{1}{x} B \varphi)| = |(B\phi, \frac{1}{x} B\chi\varphi)| \leq |(\tilde{\phi}, \frac{1}{x} \chi\tilde{\varphi})| + |(\tilde{\phi}, \frac{1}{x} [B, \chi] \varphi)| \tag{4.3}$$

where we have abbreviated  $\tilde{\phi} := B\phi$  and  $\tilde{\varphi} := B\varphi$ . Recalling that  $\chi(\frac{\mathbf{x}}{R})$  is nonvanishing only if  $x \geq R/2$  and  $\text{ran } \chi \in [0, 1]$ , the first contribution in (4.3) is easily estimated by

$$\begin{aligned}
|(\tilde{\phi}, \frac{1}{x} \chi\tilde{\varphi})| &\leq \int_{\mathbb{R}^3} d\mathbf{x} |\tilde{\phi}(\mathbf{x})| \frac{1}{x} \chi(\frac{\mathbf{x}}{R}) |\tilde{\varphi}(\mathbf{x})| \\
&\leq \frac{2}{R} \|\tilde{\phi}\| \|\tilde{\varphi}\| \leq \frac{c}{R} \|\phi\| \|\varphi\|
\end{aligned} \tag{4.4}$$

where  $\|\tilde{\phi}\| \leq \|B\| \|\phi\| \leq c_1 \|\phi\|$  was used ( $c := 2c_1^2$ ). In the second term we set  $\chi = 1 - \chi_0$  and note that  $\frac{1}{x} [B, \chi] = -\frac{1}{px} \cdot p [B, \chi_0]$  with  $\frac{1}{px}$  a bounded operator [12, p.307]. Transforming into Fourier space, we get with (3.6)

$$\begin{aligned}
\left(p \widehat{[B, \chi_0] \varphi}\right)(\mathbf{p}) &= \frac{1}{(2\pi)^{\frac{3}{2}}} p \int_{\mathbb{R}^3} d\mathbf{p}' R^3 \hat{\chi}_0((\mathbf{p} - \mathbf{p}')R) (B(\mathbf{p}) - B(\mathbf{p}')) \hat{\varphi}(\mathbf{p}') \\
&=: \int_{\mathbb{R}^3} d\mathbf{p}' k_{p[B, \chi_0]}(\mathbf{p}, \mathbf{p}') \hat{\varphi}(\mathbf{p}').
\end{aligned} \tag{4.5}$$

We define  $\psi := \frac{1}{px} \tilde{\phi}$  and apply the Lieb and Yau formula (3.1) with  $l = 1$ ,  $\boldsymbol{\xi} := \mathbf{p}$  and  $f = 1$ . Using our kernel from (4.5) we can estimate with (4.1) and (4.2)

$$\begin{aligned} I(\mathbf{p}) &:= \int_{\mathbb{R}^3} d\mathbf{p}' \frac{R^3}{(2\pi)^{\frac{3}{2}}} p |\hat{\chi}_0((\mathbf{p} - \mathbf{p}')R) | \mathbf{p} - \mathbf{p}' | |\nabla_{\mathbf{p}} B(\boldsymbol{\xi}_0)| \\ &\leq \frac{c_0}{(2\pi)^{\frac{3}{2}}} \int_{\mathbb{R}^3} d\mathbf{y} |\hat{\chi}_0(\mathbf{y})| y \frac{1}{R} \cdot \frac{p}{1 + |\mathbf{p} - (1 - \lambda)\mathbf{y}/R|}. \end{aligned} \quad (4.6)$$

The last factor is bounded for all  $p \geq 0$ , and the integral is finite because  $\hat{\chi}_0 \in \mathcal{S}(\mathbb{R}^3)$ . Thus  $I(\mathbf{p})$  is  $1/R$ -bounded, i.e.  $I(\mathbf{p}) \leq \frac{c}{R}$ .

The second integral  $J(\mathbf{p}')$  in the Lieb and Yau formula can be estimated in the same way. There, the last factor in (4.6) is replaced by  $|\mathbf{p}' + \mathbf{y}/R|/(1 + |\mathbf{p}' + \lambda\mathbf{y}/R|)$  which is also bounded for all  $p' \geq 0$ . (In the limiting case  $m = 0$ , one has  $B(\mathbf{p}) := A(p) = \frac{1}{\sqrt{2}}$  which commutes with  $\chi$ , and for  $B(\mathbf{p}) := \boldsymbol{\sigma}\mathbf{p}/p$ , one should use the explicit result  $B(\mathbf{p}) - B(\mathbf{p}') = \frac{\mathbf{p} - \mathbf{p}'}{p} (\boldsymbol{\sigma} - (\boldsymbol{\sigma}\mathbf{p}'/p')(\mathbf{p} + \mathbf{p}')/(p + p'))$  such that the last factor in (4.6) is not present.)

Together with (3.3), this proves

$$|(\tilde{\phi}, \frac{1}{x} [B, \chi] \varphi)| \leq \frac{c}{R} \|\frac{1}{px} B \phi\| \|\varphi\| \leq \frac{\tilde{c}}{R} \|\phi\| \|\varphi\|, \quad (4.7)$$

and thus the assertion of Lemma 5.

## 5. PROOF OF LEMMA 6

We have to show that  $|(\varphi, v^{(12)}\psi_n)| \leq \frac{c}{n} \|\varphi\| \|\psi_n\|$  for  $\psi_n = \varphi_n^{(1)} \phi_N^{(2)}$  with  $\varphi_n^{(1)} \in C_0^\infty(\mathbb{R}^3 \setminus B_n(0)) \otimes \mathbb{C}^2$  and  $\phi_N^{(2)} \in C_0^\infty(\mathbb{R}^3) \otimes \mathbb{C}^2$ .

For the definition of the auxiliary function  $\chi$  we note that  $\exists R_0 : \text{supp } \psi_n \subset \mathbb{R}^3 \setminus B_n(0) \cap B_{R_0}(0)$ . Choose  $n$  so large that  $R_0 < n/2$ . Then, on  $\text{supp } \psi_n : |\mathbf{x}_1 - \mathbf{x}_2| \geq x_1 - x_2 > n - \frac{n}{2} = n/2$ . Define

$$\chi_{12}\left(\frac{\mathbf{x}_1 - \mathbf{x}_2}{n}\right) := \begin{cases} 0, & |x_1 - x_2| < n/4 \\ 1, & |x_1 - x_2| \geq n/2 \end{cases}, \quad (5.1)$$

a smooth function mapping to  $[0, 1]$  with the property  $\psi_n \chi_{12} = \psi_n$ .

The operator  $v^{(12)}$ , defined in (1.2) - (1.5), reads explicitly (note that only terms even in  $\boldsymbol{\alpha}^{(k)}$  survive the projection by  $(1 + \beta^{(k)})/2$ )

$$\begin{aligned} v^{(12)} &= A(p_1)A(p_2) \frac{e^2}{|\mathbf{x}_1 - \mathbf{x}_2|} A(p_1)A(p_2) \\ &+ A(p_1)g(p_2) \boldsymbol{\sigma}^{(2)} \mathbf{p}_2 \frac{e^2}{|\mathbf{x}_1 - \mathbf{x}_2|} A(p_1)g(p_2) \boldsymbol{\sigma}^{(2)} \mathbf{p}_2 \\ &+ A(p_2)g(p_1) \boldsymbol{\sigma}^{(1)} \mathbf{p}_1 \frac{e^2}{|\mathbf{x}_1 - \mathbf{x}_2|} g(p_1) \boldsymbol{\sigma}^{(1)} \mathbf{p}_1 A(p_2) \\ &+ g(p_1) \boldsymbol{\sigma}^{(1)} \mathbf{p}_1 g(p_2) \boldsymbol{\sigma}^{(2)} \mathbf{p}_2 \frac{e^2}{|\mathbf{x}_1 - \mathbf{x}_2|} g(p_1) \boldsymbol{\sigma}^{(1)} \mathbf{p}_1 g(p_2) \boldsymbol{\sigma}^{(2)} \mathbf{p}_2 \end{aligned} \quad (5.2)$$

with  $A$  and  $g$  as in (1.3). For the present proof, we again need only the structure of each term in  $v^{(12)}$ ,  $B(\mathbf{p}_1, \mathbf{p}_2) \frac{e^2}{|\mathbf{x}_1 - \mathbf{x}_2|} B(\mathbf{p}_1, \mathbf{p}_2)$  with  $B(\mathbf{p}_1, \mathbf{p}_2) = B_1(\mathbf{p}_1) \cdot B_2(\mathbf{p}_2)$  an analytic (for  $m \neq 0$ ), bounded multiplication operator in momentum space. As in the previous proof (cf. (4.3)) we decompose

$$|(\varphi, B \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} B \psi_n)| \quad (5.3)$$

$$\leq |(B\varphi, \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} \chi_{12} B \psi_n)| + |(B\varphi, \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2| p_k} \cdot p_k [B, \chi_{12}] \psi_n)|$$

with  $k \in \{1, 2\}$ . We have  $|\mathbf{x}_1 - \mathbf{x}_2|^{-1} \leq 4/n$  on  $\text{supp } \chi_{12}$  such that the first summand in (5.3) can be estimated by  $\frac{4}{n} \|B\varphi\| \|B\psi_n\| \leq \frac{c}{n} \|\varphi\| \|\psi_n\|$ .

Consider now the second summand. The boundedness of  $(|\mathbf{x}_1 - \mathbf{x}_2| p_k)^{-1}$  is readily seen by considering e.g.  $k = 1$  and defining  $\varphi_{x_2}(\mathbf{y}_1) := \psi(\mathbf{y}_1 + \mathbf{x}_2, \mathbf{x}_2)$  for  $\psi \in \mathcal{A}(L_2(\mathbb{R}^3) \otimes \mathbb{C}^2)^2$ . Keeping  $\mathbf{x}_2$  fixed, we have with  $p_1 = \sqrt{-\nabla_{\mathbf{x}_1}^2}$  and the substitution  $\mathbf{y}_1 := \mathbf{x}_1 - \mathbf{x}_2$ ,

$$\begin{aligned} \left| \int_{\mathbb{R}^3} d\mathbf{x}_1 \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} \overline{\psi(\mathbf{x}_1, \mathbf{x}_2)} \frac{1}{p_1} \psi(\mathbf{x}_1, \mathbf{x}_2) \right| &= \left| \int_{\mathbb{R}^3} d\mathbf{y}_1 \frac{1}{y_1} \overline{\varphi_{x_2}(\mathbf{y}_1)} \frac{1}{\sqrt{-\nabla_{\mathbf{y}_1}^2}} \varphi_{x_2}(\mathbf{y}_1) \right| \\ &\leq c \int_{\mathbb{R}^3} d\mathbf{y}_1 |\varphi_{x_2}(\mathbf{y}_1)|^2 \end{aligned} \quad (5.4)$$

according to Kato's inequality. Hence,  $|(\psi, \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2| p_1} \psi)| \leq c \int_{\mathbb{R}^6} d\mathbf{x}_2 d\mathbf{y}_1 |\varphi_{x_2}(\mathbf{y}_1)|^2 = c \|\psi\|^2$ .

Furtheron, we aim at a reduction of the commutator to the one-particle case such that the proof of Lemma 5 can be mimicked.

First we note that  $\chi_{12}$  depends only on the difference of variables such that, defining  $\chi_{12,0} := 1 - \chi_{12}$ ,  $\hat{\chi}_{12,0}$  splits off a Dirac  $\delta$ -function. Using  $l = 2$  we get upon substitution of  $\mathbf{z}'_1 := \mathbf{z}_1 - \mathbf{z}_2$  for  $\mathbf{z}_1$  from the second line of (3.6)

$$\left( \widehat{\chi_{12,0} \left( \frac{\circ - *}{n} \right)} \right) (\mathbf{p}_1, \mathbf{p}_2) = \frac{1}{(2\pi)^3} n^6 \int_{\mathbb{R}^6} d\mathbf{z}_1 d\mathbf{z}_2 e^{-i\mathbf{p}_1 n \mathbf{z}_1} e^{-i\mathbf{p}_2 n \mathbf{z}_2} \chi_{12,0}(\mathbf{z}_1 - \mathbf{z}_2) \quad (5.5)$$

$$(a) = n^6 \int_{\mathbb{R}^3} d\mathbf{z}'_1 e^{-i\mathbf{p}_1 n \mathbf{z}'_1} \chi_{12,0}(\mathbf{z}'_1) \delta((\mathbf{p}_1 + \mathbf{p}_2)n) = n^3 (2\pi)^{\frac{3}{2}} \hat{\chi}_{12,0}(\mathbf{p}_1 n) \delta(\mathbf{p}_1 + \mathbf{p}_2)$$

$$(b) = n^3 (2\pi)^{\frac{3}{2}} \hat{\chi}_{12,0}(-\mathbf{p}_2 n) \delta(\mathbf{p}_1 + \mathbf{p}_2)$$

where line (b) is obtained from  $\mathbf{p}_1 = -\mathbf{p}_2$ .

Now we make use of the factorization of  $B(\mathbf{p}_1, \mathbf{p}_2)$  to write

$$p_k [B_1(\mathbf{p}_1) B_2(\mathbf{p}_2), \chi_{12}] = -p_k [B_1(\mathbf{p}_1), \chi_{12,0}] B_2 - B_1 p_k [B_2(\mathbf{p}_2), \chi_{12,0}] \quad (5.6)$$

such that the second summand in (5.3) can be split (via the triangle inequality) into two parts for each of which  $p_k$  is chosen independently ( $k = 1$  for the first and

$k = 2$  for the second term in (5.6)). Consider the kernel of the first part (cf. (4.5))

$$\begin{aligned} & k_{p_1[B_1, \chi_{12,0}]}(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}'_1, \mathbf{p}'_2) \\ &= p_1 \frac{n^3}{(2\pi)^{\frac{3}{2}}} \hat{\chi}_{12,0}((\mathbf{p}_1 - \mathbf{p}'_1)n) \delta(\mathbf{p}_1 - \mathbf{p}'_1 + \mathbf{p}_2 - \mathbf{p}'_2) (B_1(\mathbf{p}_1) - B_1(\mathbf{p}'_1)) \end{aligned} \quad (5.7)$$

where (a) is used for the Fourier transform of  $\chi_{12,0}$ . Insertion into the Lieb and Yau formula (for  $l = 2$ ) with  $f = 1$  gives

$$\begin{aligned} I(\mathbf{p}_1, \mathbf{p}_2) &:= \int_{\mathbb{R}^6} d\mathbf{p}'_1 d\mathbf{p}'_2 |k_{p_1[B_1, \chi_{12,0}]}(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}'_1, \mathbf{p}'_2)| \\ &= p_1 \frac{n^3}{(2\pi)^{\frac{3}{2}}} \int_{\mathbb{R}^3} d\mathbf{p}'_1 |\hat{\chi}_{12,0}((\mathbf{p}_1 - \mathbf{p}'_1)n)| |B_1(\mathbf{p}_1) - B_1(\mathbf{p}'_1)| \end{aligned} \quad (5.8)$$

which is independent of  $\mathbf{p}_2$  and has the identical form of (4.6) (note that the operators  $B_1$  and  $B_2$  are the same as occurred in  $b_{1m}^{(1)}$ , and  $\hat{\chi}_{12,0} \in \mathcal{S}(\mathbb{R}^3)$ ). Therefore,  $I$  (and also  $J$ ) is  $\frac{1}{n}$ -bounded by the proof of Lemma 5. The same holds true for the second part of the second summand in (5.3) which corresponds to the second term in (5.6). There, expression (b) in (5.5) has to be used. Thus, collecting results,

$$\begin{aligned} |(B\varphi, \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} [B, \chi_{12}] \psi_n)| &\leq |(\frac{1}{p_1 |\mathbf{x}_1 - \mathbf{x}_2|} B\varphi, p_1 [B_1, \chi_{12,0}] B_2 \psi_n)| \\ &\quad + |(\frac{1}{p_2 |\mathbf{x}_1 - \mathbf{x}_2|} B\varphi, B_1 p_2 [B_2, \chi_{12,0}] \psi_n)| \\ &\leq \frac{c}{n} \|\frac{1}{p_1 |\mathbf{x}_1 - \mathbf{x}_2|} B\varphi\| \|B_2 \psi_n\| + \frac{c}{n} \|B_1 \frac{1}{p_2 |\mathbf{x}_1 - \mathbf{x}_2|} B\varphi\| \|\psi_n\| \leq \frac{\tilde{c}}{n} \|\varphi\| \|\psi_n\| \end{aligned} \quad (5.9)$$

which completes the proof of Lemma 6.

## 6. PROOF OF LEMMA 3

For  $j = 0, 1$  and  $\varphi \in \mathcal{A}(C_0^\infty(\mathbb{R}^6 \setminus B_R(0)) \otimes \mathbb{C}^4)$  we have to show  $|(\phi_j \varphi, r_j \phi_j \varphi)| \leq \frac{c}{R} \|\varphi\|^2$ . (The proof for  $j = 2$  follows from the symmetry upon electron exchange.) The same strategy is used as in the previous proofs.

a)  $j = 1$ :  $r_1 = b_{1m}^{(1)} + v^{(12)}$

For the definition of the auxiliary function  $\chi$  we recall that  $\text{supp } \phi_1 \varphi \subset \mathbb{R}^6 \setminus B_R(0) \cap \{\mathbf{x} \in \mathbb{R}^6 : x_1 \geq Cx\}$ . Thus  $x = \sqrt{x_1^2 + x_2^2} \geq R$  and  $x_1 \geq CR$ . For the estimate of  $b_{1m}^{(1)}$ , we take

$$\chi_1\left(\frac{\mathbf{x}_1}{R}\right) := \begin{cases} 0, & x_1 < CR/2 \\ 1, & x_1 \geq CR \end{cases} \quad (6.1)$$

such that  $\phi_1 \varphi \chi_1 = \phi_1 \varphi$ , and we introduce  $\chi_{1,0} := 1 - \chi_1 \in \mathcal{S}(\mathbb{R}^3)$  as before. Although we are dealing here with two-particle functions, all operators ( $b_{1m}^{(1)}$  and

$\chi_1$ ) act only on particle 1. This reduces the Lieb and Yau formula to the single-particle case,

$$\begin{aligned} |(\psi, A\varphi)| &= \left| \int_{\mathbb{R}^6} d\mathbf{p}_1 d\mathbf{p}_2 \overline{\hat{\psi}(\mathbf{p}_1, \mathbf{p}_2)} \int_{\mathbb{R}^3} d\mathbf{p}'_1 k_A(\mathbf{p}_1, \mathbf{p}'_1) \hat{\varphi}(\mathbf{p}'_1, \mathbf{p}_2) \right| \\ &\leq \left( \int_{\mathbb{R}^6} d\mathbf{p}_1 d\mathbf{p}_2 |\hat{\psi}(\mathbf{p}_1, \mathbf{p}_2)|^2 I(\mathbf{p}_1) \right)^{\frac{1}{2}} \left( \int_{\mathbb{R}^6} d\mathbf{p}'_1 d\mathbf{p}_2 |\hat{\varphi}(\mathbf{p}'_1, \mathbf{p}_2)|^2 J(\mathbf{p}'_1) \right)^{\frac{1}{2}} \end{aligned} \quad (6.2)$$

where  $I$  and  $J$  are given in (3.2) with  $\boldsymbol{\xi} := \mathbf{p}_1$ . Therefore the proof of Lemma 5 can be copied to obtain

$$|(\phi_1 \varphi, b_{1m}^{(1)} \phi_1 \varphi)| \leq \frac{\tilde{c}}{R} \|\phi_1 \varphi\|^2 \leq \frac{\tilde{c}}{R} \|\varphi\|^2. \quad (6.3)$$

For the estimate of  $v^{(12)}$ , we define  $\chi$  in analogy to (5.1) by noting that additionally,  $\text{supp } \phi_1 \varphi \subset \mathbb{R}^6 \setminus B_R(0) \cap \{\mathbf{x} \in \mathbb{R}^6 : |\mathbf{x}_1 - \mathbf{x}_2| \geq Cx\}$ . Therefore,

$$\chi_{12}\left(\frac{\mathbf{x}_1 - \mathbf{x}_2}{R}\right) := \begin{cases} 0, & |\mathbf{x}_1 - \mathbf{x}_2| < CR/2 \\ 1, & |\mathbf{x}_1 - \mathbf{x}_2| \geq CR \end{cases} \quad (6.4)$$

such that again,  $\phi_1 \varphi \chi_{12} = \phi_1 \varphi$  and  $\chi_{12,0} := 1 - \chi_{12} \in \mathcal{S}(\mathbb{R}^3)$ . This enables us to adopt the proof of Lemma 6 (with  $n$  replaced by  $R$ ) to obtain

$$|(\phi_1 \varphi, v^{(12)} \phi_1 \varphi)| \leq \frac{c}{R} \|\phi_1 \varphi\|^2 \leq \frac{c}{R} \|\varphi\|^2. \quad (6.5)$$

b)  $j = 0$ :  $r_0 = b_{1m}^{(1)} + b_{1m}^{(2)}$

In this case, the support of  $\phi_j \varphi$  obeys  $\text{supp } \phi_0 \varphi \subset \mathbb{R}^6 \setminus B_R(0) \cap \{\mathbf{x} \in \mathbb{R}^6 : x_1 \geq Cx \text{ and } x_2 \geq Cx\}$ . For the discussion of  $b_{1m}^{(1)}$ , we define  $\chi_1(\mathbf{x}_1/R)$  as in (6.1) and copy the corresponding proof from a). For  $b_{1m}^{(2)}$ , we choose  $\chi_2(\mathbf{x}_2/R)$  according to (6.1) with  $x_1$  replaced by  $x_2$ . The proof is done along the same lines as for  $b_{1m}^{(1)}$ . Hence we obtain

$$\begin{aligned} |(\phi_0 \varphi, r_0 \phi_0 \varphi)| &\leq |(\phi_0 \varphi, b_{1m}^{(1)} \chi_1 \phi_0 \varphi)| + |(\phi_0 \varphi, b_{1m}^{(2)} \chi_2 \phi_0 \varphi)| \\ &\leq \frac{2c_1}{R} \|\phi_0 \varphi\|^2 \leq \frac{2c_1}{R} \|\varphi\|^2. \end{aligned} \quad (6.6)$$

## 7. PROOF OF LEMMA 1

Assume we have a normalized sequence of functions  $(\varphi_n)_{n \in \mathbb{N}}$  localized outside  $B_n(0)$  with the property (2.4) for  $h^{BR}$  and the  $\lambda \in \mathbb{R}$  under consideration. Since the normalization constant of  $\varphi_n$  tends to zero as  $n \rightarrow \infty$ , for any  $\phi \in C_0^\infty(\mathbb{R}^6)$  we have  $|(\phi, \varphi_n)| \rightarrow 0$  as  $n \rightarrow \infty$ , i.e.  $\varphi_n \xrightarrow{w} 0$ . By the Weyl criterion it follows that  $\lambda \in \sigma_{ess}(h^{BR})$ .

Conversely, let  $\lambda \in \sigma_{ess}(h^{BR})$ . Then there exists a Weyl sequence  $\psi_n \in \mathcal{A}(C_0^\infty(\mathbb{R}^3) \otimes \mathbb{C}^2)^2$ ,  $\|\psi_n\| = 1$ , with  $\psi_n \xrightarrow{w} 0$  and  $\|(h^{BR} - \lambda)\psi_n\| \rightarrow 0$  as  $n \rightarrow \infty$ .

Define a smooth symmetric (with respect to interchange of  $\mathbf{x}_1$  and  $\mathbf{x}_2$ ) function  $\chi_0 \in C_0^\infty(\mathbb{R}^6)$  mapping to  $[0, 1]$  by means of

$$\chi_0\left(\frac{\mathbf{x}}{n}\right) = \begin{cases} 1, & x \leq n \\ 0, & x > 2n \end{cases} \quad (7.1)$$

and let  $\chi_n(\mathbf{x}) := 1 - \chi_0(\frac{\mathbf{x}}{n})$ ,  $\chi_n \in C^\infty(\mathbb{R}^6 \setminus B_n(0))$ . Then  $\varphi_n := \psi_n \chi_n \in \mathcal{A}(C_0^\infty(\mathbb{R}^6 \setminus B_n(0)) \otimes \mathbb{C}^4)$  and we claim that a subsequence of  $(\varphi_n)_{n \in \mathbb{N}}$  satisfies the requirements of Lemma 1.

a)  $\|(h^{BR} - \lambda)\varphi_n\| \rightarrow 0$  as  $n \rightarrow \infty$  :

We decompose

$$\|(h^{BR} - \lambda)\chi_n\psi_n\| \leq \|\chi_n(h^{BR} - \lambda)\psi_n\| + \|[h^{BR}, \chi_0]\psi_n\|, \quad (7.2)$$

and use the equivalence (2.20) again. Concerning the first term in (7.2), we have for any  $\phi \in C_0^\infty(\mathbb{R}^6) \otimes \mathbb{C}^4$ ,

$$|(\phi, \chi_n(h^{BR} - \lambda)\psi_n)| \leq \|\chi_n\phi\| \|(h^{BR} - \lambda)\psi_n\| \rightarrow 0 \quad \text{as } n \rightarrow \infty \quad (7.3)$$

by assumption, since  $\|\chi_n\phi\| \leq \|\phi\| < \infty$ .

In order to treat the single-particle contribution to the second term in (7.2),  $T^{(k)}$  and  $b_{1m}^{(k)}$ , we change again to Fourier space and introduce the 6-dimensional Fourier transform of  $\chi_0$  according to (3.6) with  $l = 2$ . Then the kernel of  $[T^{(1)}, \chi_0]$  in momentum space reads

$$k_{[T^{(1)}, \chi_0]}(\mathbf{p}_1, \mathbf{p}_2, \mathbf{p}'_1, \mathbf{p}'_2) = \frac{n^6}{(2\pi)^3} \hat{\chi}_0((\mathbf{p}_1 - \mathbf{p}'_1)n, (\mathbf{p}_2 - \mathbf{p}'_2)n) (E_{p_1} - E_{p'_1}) \quad (7.4)$$

and by the mean value theorem (4.1), using  $\nabla_{\mathbf{p}} E_p = \nabla_{\mathbf{p}} \sqrt{p^2 + m^2} = \mathbf{p}/E_p$ ,

$$|E_{p_1} - E_{p'_1}| = |\mathbf{p}_1 - \mathbf{p}'_1| \left| \frac{\boldsymbol{\xi}}{E_{\boldsymbol{\xi}}} \right| \leq |\mathbf{p}_1 - \mathbf{p}'_1| \quad (7.5)$$

for all  $\boldsymbol{\xi} := \lambda\mathbf{p}_1 + (1 - \lambda)\mathbf{p}'_1$  with  $\lambda \in [0, 1]$ . For the integral  $I$  appearing in the Lieb and Yau formula (3.1) we have with  $f = 1$  and the substitution  $\mathbf{y}_k := (\mathbf{p}_k - \mathbf{p}'_k)n$ ,  $k = 1, 2$ ,

$$\begin{aligned} I(\mathbf{p}_1, \mathbf{p}_2) &:= \int_{\mathbb{R}^6} d\mathbf{p}'_1 d\mathbf{p}'_2 \frac{n^6}{(2\pi)^3} |\hat{\chi}_0((\mathbf{p}_1 - \mathbf{p}'_1)n, (\mathbf{p}_2 - \mathbf{p}'_2)n)| |E_{p_1} - E_{p'_1}| \\ &\leq \frac{1}{(2\pi)^3} \int_{\mathbb{R}^6} d\mathbf{y}_1 d\mathbf{y}_2 |\hat{\chi}_0(\mathbf{y}_1, \mathbf{y}_2)| y_1 \cdot \frac{1}{n} \leq \frac{c}{n} \end{aligned} \quad (7.6)$$

since  $\hat{\chi}_0 \in \mathcal{S}(\mathbb{R}^6)$ . In a similar way,  $J(\mathbf{p}'_1, \mathbf{p}'_2) \leq c/n$ , and hence

$$|(\phi, [T^{(1)}, \chi_0]\psi_n)| \leq \frac{c}{n} \|\phi\| \|\psi_n\| \quad (7.7)$$

for all  $\phi \in C_0^\infty(\mathbb{R}^6) \otimes \mathbb{C}^4$ . For the operator  $b_{1m}^{(1)}$  we can proceed as in the proof of Lemma 5, because (according to (7.6)) the two-particle nature of  $\chi_0$  does not affect the convergence of the single-particle integrals.

For the estimate of the remaining commutator,  $[v^{(12)}, \chi_0]$ , we follow section 5 to split it into commutators of  $\chi_0$  with single-particle (bounded) operators  $B_1(\mathbf{p}_1)$ ,  $B_2(\mathbf{p}_2)$ . The only difference as compared to the proof of Lemma 6 lies in the two-particle nature of  $\chi_0$  (cf. (7.4) in place of (5.7)), but again this does not affect the convergence of the integrals. Thus we get

$$\begin{aligned} |(\phi, [h^{BR}, \chi_0] \psi_n)| &\leq \sum_{k=1}^2 \left( |(\phi, [T^{(k)}, \chi_0] \psi_n)| + |(\phi, [b_{1m}^{(k)}, \chi_0] \psi_n)| \right) \\ &+ |(\phi, [v^{(12)}, \chi_0] \psi_n)| \leq \frac{c}{n} \|\phi\| \|\psi_n\| \end{aligned} \quad (7.8)$$

with the generic constant  $c$ .

b)  $\|\varphi_n\| \neq 0$ :

We show that for any  $\epsilon > 0$  there is an  $N_0 \in \mathbb{N}$  such that  $\|\chi_0 \psi_n\| < \epsilon$  for all  $n \geq N_0$ . Then  $\|(1 - \chi_n) \psi_n\| = \|\psi_n - \varphi_n\| < \epsilon$ . As a consequence,  $\|\varphi_n\| \neq 0$  for sufficiently large  $n > n_0 \geq N_0$  since  $\|\psi_n\| = 1$  (i.e. one can choose a subsequence of  $(\varphi_n)_{n \in \mathbb{N}}$  with normalizable elements).

Since  $h^{BR}$  is a self-adjoint positive operator,  $h^{BR} + 1 > 0$  with a bounded inverse. Then following [4] we estimate

$$\begin{aligned} \|\chi_0 \psi_n\| &= \|\chi_0 (h^{BR} + 1)^{-1} [(h^{BR} - \lambda) + (1 + \lambda)] \psi_n\| \\ &\leq \|\chi_0 (h^{BR} + 1)^{-1}\| \|(h^{BR} - \lambda) \psi_n\| + |1 + \lambda| \|\chi_0 (h^{BR} + 1)^{-1} \psi_n\|. \end{aligned} \quad (7.9)$$

Since  $\chi_0 \in C_0^\infty(\mathbb{R}^6)$  and by assumption  $\exists N_1 : \|(h^{BR} - \lambda) \psi_n\| < \tilde{\epsilon}$  for  $n \geq N_1$ , the first term is bounded by, say,  $c_1 \tilde{\epsilon}$  for  $n \geq N_1$ .

For the second term we show that the bounded function  $\chi_0$  is relatively compact with respect to  $h^{BR}$ . We write

$$\chi_0 (h^{BR} + 1)^{-1} = \chi_0 (T + 1)^{-\frac{1}{2}} (T + 1)^{\frac{1}{2}} (h^{BR} + 1)^{-1} \quad (7.10)$$

and note that  $(T + 1)^{\frac{1}{2}} (h^{BR} + 1)^{-\frac{1}{2}} \cdot (h^{BR} + 1)^{-\frac{1}{2}}$  is a product of bounded operators since  $V$  is  $T$ -from bounded (with form bound  $< 1$  for  $\gamma < \gamma_{BR}$ ).

For the compactness of  $\chi_0 (T + 1)^{-\frac{1}{2}}$  we apply a theorem ([22, p.115], [21, Lemma 6.9] stating that for bounded functions  $f(x)$  and  $g(p)$  with  $f, g : [0, \infty) \rightarrow \mathbb{C}$  and  $\lim_{x \rightarrow \infty} f(x) = 0 = \lim_{p \rightarrow \infty} g(p)$  the product  $K := gf$  is compact.

Clearly, both  $\chi_0$  and  $(T + 1)^{-\frac{1}{2}} = (\sqrt{p_1^2 + m^2} + \sqrt{p_2^2 + m^2} + 1)^{-\frac{1}{2}}$  are bounded functions, going to zero as  $x \rightarrow \infty$  and  $p := \sqrt{p_1^2 + p_2^2} \rightarrow \infty$ , respectively. Therefore,  $\chi_0 (h^{BR} + 1)^{-1}$  is compact and maps the weakly convergent sequence  $(\psi_n)_{n \in \mathbb{N}}$  into a strongly convergent sequence. So the second term in (7.9) can be estimated by  $c_2 \tilde{\epsilon}$  for, say,  $n \geq N_2$ . This proves the assertion  $\|\chi_0 \psi_n\| < \epsilon := (c_1 + c_2) \tilde{\epsilon}$  for  $n \geq N_0 := \max\{N_1, N_2\}$ .

## 8. PROOF OF LEMMA 4

We have to show that for  $\varphi \in \mathcal{A}(C_0^\infty(\mathbb{R}^6 \setminus B_R(0)) \otimes \mathbb{C}^4)$  and  $\chi(\frac{\mathbf{x}}{R})$  the auxiliary function from (3.4) with  $l = 2$ ,  $(\phi_j \varphi, [b_{1m}^{(1)}, \phi_j \chi] \varphi)$  as well as  $(\phi_j \varphi, [v^{(1,2)}, \phi_j \chi] \varphi)$  are uniformly  $1/R$ -bounded.

Let us denote  $\phi_j \chi =: \psi_j$ . We have  $\text{supp } \phi_j \chi \subseteq \text{supp } \chi$  which is located outside  $B_{R/2}(0)$  such that the scaling holds,

$$\psi_j(\mathbf{x}) = \phi_j(\mathbf{x}) \chi\left(\frac{\mathbf{x}}{R}\right) = \phi_j\left(\frac{\mathbf{x}}{R/2}\right) \chi\left(\frac{\mathbf{x}}{R}\right) \quad (8.1)$$

for  $R \geq 2$ . Since  $\phi_j$  and  $\chi$  are analytic functions in  $\text{supp } \chi$  we can apply the mean value theorem, with  $\mathbf{x} = (\mathbf{x}_1, \mathbf{x}_2)$ ,

$$|\psi_j(\mathbf{x}_1, \mathbf{x}_2) - \psi_j(\mathbf{x}'_1, \mathbf{x}_2)| = |\mathbf{x}_1 - \mathbf{x}'_1| |(\nabla_{\mathbf{x}_1} \psi_j)(\boldsymbol{\xi}, \mathbf{x}_2)| \quad (8.2)$$

with  $\boldsymbol{\xi}$  some value on the line between  $\mathbf{x}_1$  and  $\mathbf{x}'_1$ . Since from (3.5),  $\chi' = -\chi'_0 \in \mathcal{S}(\mathbb{R}^6)$  is a bounded function, as is  $\phi'_j$  (because  $\phi_j \in C^\infty(\mathbb{R}^6)$  is defined on the compact unit sphere and is homogeneous of degree zero outside the unit ball), we can estimate

$$\left| \left( \nabla_{\mathbf{x}_1} \phi_j\left(\frac{\mathbf{x}}{R/2}\right) \chi\left(\frac{\mathbf{x}}{R}\right) \right) (\boldsymbol{\xi}, \mathbf{x}_2) \right| = |\phi_j| \frac{1}{R} |\chi'(\frac{\boldsymbol{\xi}}{R}, \frac{\mathbf{x}_2}{R})| + |\chi| \frac{1}{R/2} |\phi'_j(\boldsymbol{\xi}, \mathbf{x}_2)| \leq \frac{c_0}{R} \quad (8.3)$$

(the prime referring to the derivative with respect to the first entry).  $c_0$  is a suitable constant which can be chosen independently of  $j$ .

a) Using the explicit form (1.5) of  $b_{1m}^{(1)}$ , we decompose the commutator in the following way,

$$\begin{aligned} [b_{1m}^{(1)}, \psi_j] &= -\gamma \left\{ [A, \psi_j] \frac{1}{x_1} A + [\sigma^{(1)} \mathbf{p}_1 g, \psi_j] \frac{1}{x_1} g \sigma^{(1)} \mathbf{p}_1 \right. \\ &\quad \left. + A \frac{1}{x_1} [A, \psi_j] + \sigma^{(1)} \mathbf{p}_1 g \frac{1}{x_1} [g \sigma^{(1)} \mathbf{p}_1, \psi_j] \right\}. \end{aligned} \quad (8.4)$$

Since the two terms in the second line of (8.4) are (up to a sign) the hermitean conjugate of the first line, and the interchange of  $\phi_j \varphi$  with  $\varphi$  in the quadratic form plays no role, we need not discuss these terms separately.

The terms in the first line of (8.4) have the structure  $[\mathcal{O}, \psi_j] \frac{1}{x_1} B$  with  $\mathcal{O}$  and  $B$  multiplication operators in momentum space and  $B$  bounded. Provided we can show

$$|(\varphi, [\mathcal{O}, \psi_j] \frac{1}{x_1} \psi)| \leq \frac{c}{R} \|\varphi\| \|\psi\|, \quad (8.5)$$

we can estimate  $|(\phi_j \varphi, [\mathcal{O}, \psi_j] \frac{1}{x_1} B \varphi)| \leq \frac{c}{R} \|\phi_j \varphi\| \|B \varphi\| \leq \frac{c_1}{R} \|\varphi\|^2$  and we are done. The proof of (8.5) is based on the following estimate of the kernel  $\check{k}_{\mathcal{O}}$  of  $\mathcal{O}$  in coordinate space,

$$|\check{k}_{\mathcal{O}}(\mathbf{x}_1, \mathbf{x}'_1)| \leq \frac{c}{|\mathbf{x}_1 - \mathbf{x}'_1|^3}, \quad (8.6)$$



for our two operators of interest,  $A$  and  $\sigma^{(1)}\mathbf{p}_1g$ . Assuming (8.6) to hold, we apply the coordinate-space version of the Lieb and Yau formula (3.1) to the l.h.s. of (8.5), identifying  $\boldsymbol{\xi}$  with  $(\mathbf{x}_1, \mathbf{x}_2)$ . Then we have to estimate the two integrals from (3.2),  $I$  and  $J$ . For  $I$  we have

$$I(\mathbf{x}_1, \mathbf{x}_2) := \int_{\mathbb{R}^6} d\mathbf{x}'_1 d\mathbf{x}'_2 |\check{k}_{[\mathcal{O}, \psi_j]_{\perp_1}}(\mathbf{x}_1, \mathbf{x}'_1)| \delta(\mathbf{x}_2 - \mathbf{x}'_2) \frac{f(\mathbf{x}_1, \mathbf{x}_2)}{f(\mathbf{x}'_1, \mathbf{x}'_2)} \quad (8.7)$$

with the convergence generating function chosen to be  $f(\mathbf{x}_1, \mathbf{x}_2) = x_1^{3/2}$ . The delta function appears because the momentum operator  $\mathcal{O}$  does not affect particle 2. With the help of (8.6) and the mean value theorem (8.2) and (8.3) we obtain

$$\begin{aligned} I(\mathbf{x}_1, \mathbf{x}_2) &= \int_{\mathbb{R}^3} d\mathbf{x}'_1 |\check{k}_{\mathcal{O}}(\mathbf{x}_1, \mathbf{x}'_1)| |\psi_j(\mathbf{x}_1, \mathbf{x}_2) - \psi_j(\mathbf{x}'_1, \mathbf{x}_2)| \frac{1}{x'_1} \cdot \frac{x_1^{3/2}}{x_1^{3/2}} \\ &\leq \frac{c_0 c}{R} \int_{\mathbb{R}^3} d\mathbf{x}'_1 \frac{1}{|\mathbf{x}_1 - \mathbf{x}'_1|^3} |\mathbf{x}_1 - \mathbf{x}'_1| \frac{x_1^{3/2}}{x_1^{5/2}}. \end{aligned} \quad (8.8)$$

Using spherical coordinates, the angular integration is performed by means of  $\int_{S^2} d\omega' \frac{1}{|\mathbf{x}_1 - \mathbf{x}'_1|^2} = \frac{2\pi}{x_1 x'_1} \ln \left| \frac{x_1 + x'_1}{x_1 - x'_1} \right|$  [8]. With the substitution  $x'_1 =: x_1 z'$  we get for the r.h.s. of (8.8),

$$I(\mathbf{x}_1, \mathbf{x}_2) \leq \frac{2\pi c_0 c}{R} \int_0^\infty \frac{dz'}{z'^{3/2}} \ln \left| \frac{1+z'}{1-z'} \right| \leq \frac{C}{R} \quad (8.9)$$

since the integral is convergent. For the integral  $J$  we use the identical estimates. Then with  $x_1 =: x'_1 z$ ,

$$\begin{aligned} J(\mathbf{x}'_1, \mathbf{x}'_2) &:= \int_{\mathbb{R}^6} d\mathbf{x}_1 d\mathbf{x}_2 |\check{k}_{[\mathcal{O}, \psi_j]_{\perp_1}}(\mathbf{x}_1, \mathbf{x}'_1)| \delta(\mathbf{x}_2 - \mathbf{x}'_2) \frac{x_1^{3/2}}{x_1^{3/2}} \\ &\leq \frac{2\pi c_0 c}{R} \int_0^\infty \frac{dz}{z^{1/2}} \ln \left| \frac{1+z}{1-z} \right| \leq \frac{\tilde{C}}{R}. \end{aligned} \quad (8.10)$$

From (8.9) and (8.10), the Lieb and Yau formula together with (3.3) provides the desired result (8.5).

It remains to show the estimate (8.6) for the kernel of our operators. Let us first consider the limiting case  $m = 0$  as this is very simple.

For  $m = 0$ ,  $\sigma^{(1)}\mathbf{p}_1g = \frac{1}{\sqrt{2}}\sigma^{(1)}\frac{\mathbf{p}_1}{p_1}$ , while  $A$  is a constant which need not be considered. From (3.7), using the Fourier transform of the Coulomb potential,

$$\begin{aligned} \check{k}_{\sigma^{(1)}\mathbf{p}_1g}(\mathbf{x}_1, \mathbf{x}'_1) &= -i\sigma^{(1)}\nabla_{\mathbf{x}_1} \frac{1}{\sqrt{2}} \frac{4\pi}{(2\pi)^3} \frac{1}{|\mathbf{x}_1 - \mathbf{x}'_1|^2} \\ &= \frac{i\sigma^{(1)}}{\sqrt{2}\pi^2} \frac{\mathbf{x}_1 - \mathbf{x}'_1}{|\mathbf{x}_1 - \mathbf{x}'_1|^4}, \end{aligned} \quad (8.11)$$

such that the desired estimate (8.6) follows immediately.

For  $m \neq 0$ , we have to consider the two functions  $g(p_1)$  and  $A(p_1)$  defined in (1.3). Without loss of generality we can set  $m = 1$  (otherwise, due to scaling, one would have to consider  $m\tilde{x}$  in place of  $\tilde{x}$  in the integrand of (3.7)). Then

$$\begin{aligned} g(p_1) &= \frac{1}{\sqrt{2}} \left( \frac{1}{p_1^2 + 1 + \sqrt{p_1^2 + 1}} \right)^{\frac{1}{2}} \\ A(p_1) &= \frac{1}{\sqrt{2}} \left( 1 + \frac{1}{\sqrt{p_1^2 + 1}} \right)^{\frac{1}{2}} = \frac{1}{\sqrt{2}} + \tilde{g}(p_1), \\ \tilde{g}(p_1) &:= \frac{1}{\sqrt{2}} \frac{1}{\sqrt{p_1^2 + 1} + \sqrt{p_1^2 + 1 + \sqrt{p_1^2 + 1}}} \end{aligned} \quad (8.12)$$

where  $\tilde{g}(p_1) \sim 2^{-\frac{3}{2}}/p_1$  for  $p_1 \rightarrow \infty$ .

Both functions, if extended to the complex plane, have branch points at  $p_1 = \pm i$  and are analytic in the strip  $\{z \in \mathbb{C} : |\operatorname{Im} z| < 1\}$  if the cuts are chosen from  $i$  to  $\infty$  and from  $-\infty$  to  $-i$ , respectively. According to a corollary of the Paley-Wiener theorem [17, Thm IX.14], for  $\tilde{x} := |\mathbf{x}'_1 - \mathbf{x}_1| > 0$ , it follows that  $\check{k}_{\tilde{g}}$  and  $\check{k}_g$  (if defined with convergence generating factors) are bounded continuous functions decaying exponentially at  $\tilde{x} \rightarrow \infty$ , i.e. for any  $b < 1$  and any  $\delta > 0$ , there exists a constant  $c_1 > 0$ :

$$|\check{k}_{\tilde{g}}(\tilde{x})| \leq c_1 e^{-b\tilde{x}} \quad \text{for } \tilde{x} \geq \delta, \quad (8.13)$$

and similarly for  $\check{k}_g$ . Here and in the following we have to introduce the convergence generating factors by means of replacing for  $\epsilon > 0$  the integral in (3.7) by

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\tilde{x}} \int_0^\infty p_1 dp_1 \sin p_1 \tilde{x} \tilde{g}(p_1) e^{-\epsilon p_1}. \quad (8.14)$$

In determining the behaviour of  $\check{k}_{\tilde{g}}$  near  $\tilde{x} = 0$ , we apply two partial integrations to (8.14) and obtain

$$\begin{aligned} \frac{1}{\tilde{x}} \left| \int_0^\infty dp_1 \sin p_1 \tilde{x} (p_1 \tilde{g} e^{-\epsilon p_1}) \right| &= \frac{1}{\tilde{x}^3} \left| \int_0^\infty dp_1 \sin p_1 \tilde{x} \left\{ \left( \frac{d^2}{dp_1^2} p_1 \tilde{g} \right) e^{-\epsilon p_1} \right. \right. \\ &\quad \left. \left. - 2 \frac{1}{p_1} \left( \frac{d}{dp_1} p_1 \tilde{g} \right) \cdot [\epsilon p_1 e^{-\epsilon p_1}] + \frac{\tilde{g}}{p_1} [(\epsilon p_1)^2 e^{-\epsilon p_1}] \right\} \right| \leq \frac{c}{\tilde{x}^3} \end{aligned} \quad (8.15)$$

because the boundary terms vanish,  $|\sin p_1 \tilde{x}| \leq 1$  and the term in curly brackets is bounded by  $\frac{c}{(p_1+1)^2}$  for all  $\epsilon$ , which is an integrable function.

The continuity of  $\check{k}_{\tilde{g}}$  at  $\tilde{x} > 0$  together with (8.13) thus ensures the estimate  $|\check{k}_{\tilde{g}}(\tilde{x})| \leq \frac{c}{\tilde{x}^3}$ .

For the second kernel,  $\check{k}_{\sigma^{(1)} \mathbf{p}_1 g}(\tilde{x})$ , we need more careful estimates due to the presence of the derivative  $\sigma^{(1)} \nabla_{\tilde{\mathbf{x}}}$  in (3.7). First we prove analyticity of the kernel

$\check{k}_g$ . We use (8.14) with  $\tilde{g}$  replaced by  $g$ . Writing  $\sin p_1 \tilde{x} = \frac{1}{2i}(e^{ip_1 \tilde{x}} - e^{-ip_1 \tilde{x}})$ , substituting  $p_1 = iy$  and using symmetry, (8.14) turns into

$$\lim_{\epsilon \rightarrow 0} \frac{1}{2i\tilde{x}} \left( \int_{-\infty}^0 \tilde{f}(y) e^{-i\epsilon y} dy + \int_0^{i\infty} \tilde{f}(y) e^{i\epsilon y} dy \right)$$

$$\tilde{f}(y) := y e^{-\tilde{x}y} \frac{1}{\sqrt{2}} \left( \frac{1}{1 - y^2 + \sqrt{1 - y^2}} \right)^{\frac{1}{2}} \quad (8.16)$$

Using analyticity of the integrands outside the cuts as well as their exponential decay as  $\text{Re}(y) \rightarrow \infty$ , we can use Cauchy's integral theorem to deform the integration paths to the real axis. The resulting integrals are finite for all  $\tilde{x} > 0$  and  $\epsilon = 0$  since the singularity at  $y = 1$  is integrable. Thus (8.16) gives

$$\frac{1}{\tilde{x}\sqrt{2}} \int_1^{\infty} y dy e^{-\tilde{x}y} \text{Re} \left( \frac{1}{y^2 - 1 + i\sqrt{y^2 - 1}} \right)^{\frac{1}{2}} \quad (8.17)$$

with positive choice of the real part. Its derivative with respect to  $\tilde{x}$  also converges absolutely and hence (8.17) represents an analytic function for  $\tilde{x} > 0$ .

For the large- $\tilde{x}$  behaviour we note that an expansion around the branch point  $p_1 = i$  provides, as outlined by Murray [15],

$$\frac{1}{\tilde{x}} \int_0^{\infty} p_1 dp_1 \sin p_1 \tilde{x} g(p_1) e^{-\epsilon p_1^4} \sim z_0 \frac{e^{-\tilde{x}}}{\tilde{x}^{7/4}} e^{-\epsilon} \quad \text{as } \tilde{x} \rightarrow \infty \quad (8.18)$$

with a constant  $z_0$ .

For the estimate of  $\check{k}_g$  near  $\tilde{x} = 0$ , we separate the 'Coulombic' tail from  $g$ ,  $g(p) = \frac{1}{\sqrt{2}p} + (g(p) - \frac{1}{\sqrt{2}p})$  and obtain with (8.11) and (3.7)

$$\check{k}_g(\tilde{x}) = \frac{1}{2\sqrt{2}\pi^2} \left( \frac{1}{\tilde{x}^2} - \lim_{\epsilon \rightarrow 0} I_{\epsilon}(\tilde{x}) \right) \quad (8.19)$$

$$I_{\epsilon}(\tilde{x}) := \frac{1}{\tilde{x}} \int_0^{\infty} dp_1 \sin p_1 \tilde{x} \frac{(1 + \sqrt{p_1^2 + 1})^{\frac{1}{2}}}{(p_1^2 + 1)^{\frac{1}{4}} [p_1 + (p_1^2 + 1 + \sqrt{p_1^2 + 1})^{\frac{1}{2}}]} e^{-\epsilon p_1}.$$

The fraction multiplying  $\sin p_1 \tilde{x}$  decreases according to  $1/p_1$  for  $p_1 \rightarrow \infty$ . With one partial integration, we get

$$I_{\epsilon}(\tilde{x}) = \frac{1}{\tilde{x}^2} + \frac{1}{\tilde{x}^2} \int_0^{\infty} dp_1 \cos p_1 \tilde{x} \left\{ \left( \frac{d}{dp_1} [\dots] \right) e^{-\epsilon p_1} - \frac{[\dots]}{p_1} (\epsilon p_1 e^{-\epsilon p_1}) \right\} \quad (8.20)$$

where the first term comes from the boundary at  $p_1 = 0$  and  $[\dots]$  denotes the fraction in (8.19). The integral is finite (and independent of  $\tilde{x}$  for  $\tilde{x} \rightarrow 0$ ) because the curly bracket can be estimated by  $\frac{c_0}{(1+p_1)^2}$  (independent of  $\epsilon$ ) which is integrable. Hence,  $\check{k}_g(\tilde{x}) \sim \frac{\tilde{c}}{\tilde{x}^2}$  for  $\tilde{x} \rightarrow 0$  with some constant  $\tilde{c}$ . Performing the derivative inherent in  $\sigma^{(1)} \mathbf{p}_1$ , we find  $\check{k}_{\sigma^{(1)} \mathbf{p}_1 g}(\tilde{x}) \sim 2i\tilde{c} \frac{\sigma^{(1)} \tilde{x}}{\tilde{x}^4}$  ( $\tilde{x} \rightarrow 0$ ) and

$\sim iz_0 \frac{\sigma^{(1)} \tilde{x}}{\tilde{x}^4} (\tilde{x}^{5/4} + \frac{7}{4} \tilde{x}^{1/4}) e^{-\tilde{x}}$  for  $\tilde{x} \rightarrow \infty$ . From the analyticity of  $\check{k}_g$  we finally obtain

$$|\check{k}_{\sigma^{(1)} \mathbf{p}_1 g}(\tilde{x})| \leq \frac{c}{\tilde{x}^3} \quad \text{for all } \tilde{x} > 0. \quad (8.21)$$

b) Let us now turn to the commutator with the two-body interaction  $v^{(12)}$ . According to the explicit expression (5.2) for  $v^{(12)}$ , we split  $[v^{(12)}, \psi_j] =: e^2 \sum_{k=1}^4 M_k$  into four contributions and write them in the following way (with the short-hand notation  $A_k := A(p_k)$ ,  $g_k := g(p_k)$ ,  $k = 1, 2$ ).

$$\begin{aligned} M_1 &:= [A_1, \psi_j] \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} \cdot |\mathbf{x}_1 - \mathbf{x}_2| A_2 \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} A_1 A_2 \\ &\quad + A_1 [A_2, \psi_j] \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} A_1 A_2 - h.c. \\ M_2 &:= [A_1, \psi_j] \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} \cdot |\mathbf{x}_1 - \mathbf{x}_2| g_2 \sigma^{(2)} \mathbf{p}_2 \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} A_1 g_2 \sigma^{(2)} \mathbf{p}_2 \\ &\quad + A_1 [g_2 \sigma^{(2)} \mathbf{p}_2, \psi_j] \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} A_1 g_2 \sigma^{(2)} \mathbf{p}_2 - h.c. \\ M_3 &:= [g_1 \sigma^{(1)} \mathbf{p}_1, \psi_j] \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} \cdot |\mathbf{x}_1 - \mathbf{x}_2| g_2 \sigma^{(2)} \mathbf{p}_2 \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} g_1 \sigma^{(1)} \mathbf{p}_1 g_2 \sigma^{(2)} \mathbf{p}_2 \\ &\quad + g_1 \sigma^{(1)} \mathbf{p}_1 [g_2 \sigma^{(2)} \mathbf{p}_2, \psi_j] \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} g_1 \sigma^{(1)} \mathbf{p}_1 g_2 \sigma^{(2)} \mathbf{p}_2 - h.c. \end{aligned} \quad (8.22)$$

where  $h.c.$  denotes the hermitean conjugate, and  $M_3$  results from  $M_2$  upon interchanging  $\mathbf{p}_1, \sigma^{(1)}$  with  $\mathbf{p}_2, \sigma^{(2)}$  and therefore need not be considered separately.

In addition to estimate the commutators  $[\mathcal{O}, \psi_j] \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|}$  (with the same operators  $\mathcal{O}$  as in  $a$ ) except for a possible particle exchange), we have to prove boundedness of  $|\mathbf{x}_1 - \mathbf{x}_2| \mathcal{O} \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|}$  which appears as a factor in the first contribution to  $M_k$ ,  $k = 1, \dots, 4$ .

For the estimate of the commutators we proceed as in  $a$ ) except for the choice  $f(\mathbf{x}_1, \mathbf{x}_2) = |\mathbf{x}_1 - \mathbf{x}_2|^{3/2}$ . With the inequality (8.6) for the kernel of  $\mathcal{O}$  and the mean value theorem (8.2) and (8.3) we obtain for  $I$ , defined as the integral over the kernel of  $[\mathcal{O}, \psi_j] \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|}$  where  $\mathcal{O}$  depends e.g. on  $\mathbf{p}_2$ ,

$$\begin{aligned} I(\mathbf{x}_1, \mathbf{x}_2) &:= \int_{\mathbb{R}^3} d\mathbf{x}'_2 |\check{k}_{\mathcal{O}}(\mathbf{x}_2, \mathbf{x}'_2)| |\psi_j(\mathbf{x}_1, \mathbf{x}_2) - \psi_j(\mathbf{x}_1, \mathbf{x}'_2)| \frac{1}{|\mathbf{x}_1 - \mathbf{x}'_2|} \frac{f(\mathbf{x}_1, \mathbf{x}_2)}{f(\mathbf{x}_1, \mathbf{x}'_2)} \\ &\leq \frac{c_0 c}{R} \int_{\mathbb{R}^3} d\mathbf{x}'_2 \frac{1}{|\mathbf{x}_2 - \mathbf{x}'_2|^3} |\mathbf{x}_2 - \mathbf{x}'_2| \frac{1}{|\mathbf{x}_1 - \mathbf{x}'_2|} \frac{|\mathbf{x}_1 - \mathbf{x}_2|^{3/2}}{|\mathbf{x}_1 - \mathbf{x}'_2|^{3/2}}. \end{aligned} \quad (8.23)$$

We abbreviate  $\mathbf{x}_0 := \mathbf{x}_2 - \mathbf{x}_1$  and substitute  $\mathbf{x}'_2 - \mathbf{x}_1 = x_0 \mathbf{y}$ . Then the second line in (8.23) is written as ( $\mathbf{e}_{x_0} := \mathbf{x}_0/x_0$ )

$$\begin{aligned} I(\mathbf{x}_1, \mathbf{x}_2) &\leq \frac{c_0 c}{R} \int_{\mathbb{R}^3} d\mathbf{y} \frac{1}{|\mathbf{e}_{x_0} - \mathbf{y}|^2} \frac{1}{y^{5/2}} \\ &= \frac{2\pi c_0 c}{R} \int_0^\infty \frac{dy}{y^{3/2}} \ln \left| \frac{1+y}{1-y} \right| \leq \frac{C}{R} \end{aligned} \quad (8.24)$$

with the same estimate as in (8.9) above. Likewise, the estimate (8.10) holds for the second integral  $J$ .

Finally we have to show the boundedness of  $|\mathbf{x}_1 - \mathbf{x}_2| \mathcal{O} \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|}$ . Since commutators help to regularize integrals, we decompose

$$|\mathbf{x}_1 - \mathbf{x}_2| \mathcal{O} \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|} = \mathcal{O} + [|\mathbf{x}_1 - \mathbf{x}_2|, \mathcal{O}] \frac{1}{|\mathbf{x}_1 - \mathbf{x}_2|}. \quad (8.25)$$

$\mathcal{O}$  is bounded so we can concentrate on the second term. The difference to the commutators occurring in (8.22) is the replacement of  $\psi_j$  by  $|\mathbf{x}_1 - \mathbf{x}_2|$ . From the mean value theorem we get (for  $\mathcal{O}$  depending on  $\mathbf{p}_2$ )

$$||\mathbf{x}_1 - \mathbf{x}_2| - |\mathbf{x}_1 - \mathbf{x}'_2|| = \left| (\mathbf{x}_2 - \mathbf{x}'_2) \frac{\boldsymbol{\xi} - \mathbf{x}_1}{|\boldsymbol{\xi} - \mathbf{x}_1|} \right| \leq |\mathbf{x}_2 - \mathbf{x}'_2| \quad (8.26)$$

with  $\boldsymbol{\xi}$  some point between  $\mathbf{x}_2$  and  $\mathbf{x}'_2$ . This means that the earlier estimates (8.24) and (8.10) hold except for the factor  $c_0/R$  from (8.3) which is not present now. This proves the desired boundedness.

The operators in  $M_k$ ,  $k = 1, \dots, 4$  from (8.22) which are not yet considered, are all bounded. Applying the Lieb and Yau formula, this finally shows that  $|(\phi_j \varphi, [v^{(12)}, \psi_j] \varphi)| \leq \frac{c}{R} \|\varphi\|^2$  and completes the proof of the HVZ theorem.

#### ACKNOWLEDGMENT

I should like to thank H.Siedentop, H.Kalf, R.Lasser, L.Erdős, S.Vugalter and S.Morozov for stimulating discussions and critical comments. It is a pleasure to thank A.Sobolev for his hint to work in momentum space and to use the auxiliary function in the proofs, as well as for his hospitality during my stay at the University of Sussex in 2005. Support by the EU Network Analysis and Quantum (contract HPRN-CT-2002-00277) is gratefully acknowledged.

#### REFERENCES

- [1] Balinsky A.A. and Evans W.D., On the Brown-Ravenhall relativistic Hamiltonian and the stability of matter, *Stud. Adv. Math.* **16**, 1-9 (2000).
- [2] Brown G.E. and Ravenhall D.G., On the interaction of two electrons, *Proc. Roy. Soc. London* **A208**, 552-559 (1951).
- [3] Burenkov V.I. and Evans W.D., On the evaluation of the norm of an integral operator associated with the stability of one-electron atoms, *Proc. Roy. Soc. (Edinburgh)* **128A**, 993-1005 (1998).
- [4] Cycon H.L., Froese R.G., Kirsch W., Simon B., *Schrödinger Operators with Application to Quantum Mechanics and Global Geometry*, Text and Monographs in Physics, 1st Edition, Springer-Verlag, Berlin, 1987.
- [5] Douglas M. and Kroll N.M., Quantum electrodynamical corrections to the fine structure of helium, *Ann. Phys. (N.Y.)* **82**, 89-155 (1974).
- [6] Evans W.D., Perry P., Siedentop H., The spectrum of relativistic one-electron atoms according to Bethe and Salpeter, *Commun. Math. Phys.* **178**, 733-746 (1996).
- [7] Georgescu V. and Iftimovici A., Crossed products of  $C^*$ -algebras and spectral analysis of quantum Hamiltonians, *Commun. Math. Phys.* **228**, 519-560 (2002).
- [8] Gradshteyn I.S. and Ryzhik I.M., *Table of Integrals, Series and Products*, Academic Press, New York, 1965.

- [9] Herbst I.W., Spectral theory of the operator  $(p^2 + m^2)^{\frac{1}{2}} - Ze^2/r$ . *Commun. Math. Phys.* **53**, 285-294 (1977).
- [10] Hunziker W., On the spectra of Schrödinger multiparticle Hamiltonians, *Helv. Phys. Acta* **39**, 451-462 (1966).
- [11] Jakubaša-Amundsen D.H., Pseudorelativistic operator for a two-electron ion, *Phys. Rev.* **A71**, 032105, 1-8 (2005).
- [12] Kato T., *Perturbation Theory for Linear Operators*, Springer-Verlag, Berlin, 1980.
- [13] Lewis R.T., Siedentop H., Vugalter S., The essential spectrum of relativistic multi-particle operators, *Ann. Inst. Henri Poincaré* **67**, No.1, 1-28 (1997).
- [14] Lieb E.H. and Yau H.-T., The stability and instability of relativistic matter, *Commun. Math. Phys.* **118**, 177-213 (1988).
- [15] Murray J.D., *Asymptotic Analysis*, Clarendon Press, Oxford, 1974, §2, 5.
- [16] Reed M. and Simon B., *Functional Analysis*, Vol. I of Methods of Modern Mathematical Physics, Academic Press, New York, 1980.
- [17] Reed M. and Simon B., *Fourier Analysis, Self-Adjointness*, Vol. II of Methods of Modern Mathematical Physics, Academic Press, New York, 1975.
- [18] Reed M. and Simon B., *Analysis of Operators*, Vol. IV of Methods of Modern Mathematical Physics, Academic Press, New York, 1978.
- [19] Siedentop H. and Stockmeyer E., An analytic Douglas-Kroll-Hess method, *Phys. Lett.* **A341**, 473-478 (2005).
- [20] Simon B., Geometric methods in multiparticle quantum systems, *Commun. Math. Phys.* **55**, 259-274 (1977).
- [21] Teschl G., Lecture Notes on *Schrödinger Operators*, Section 9.4, 2004.  
<http://www.mat.univie.ac.at/~gerald/ftp/index.html>
- [22] Thaller B., *The Dirac equation*, Springer-Verlag, Berlin, 1992.
- [23] Tix C., Lower bound for the ground state energy of the no-pair Hamiltonian, *Phys. Lett.* **B405**, 293-296 (1997).
- [24] Tix C., Strict positivity of a relativistic Hamiltonian due to Brown and Ravenhall, *Bull. London Math. Soc.* **30**, 283-290 (1998).
- [25] Weder R.A., Spectral analysis of pseudodifferential operators, *J. Functional Analysis* **20**, 319-337 (1975).
- [26] van Winter C., Theory of finite systems of particles I. The Green function, *Mat. Fys. Dan. Vid. Selsk.* **2** No.8, 1-60 (1964).
- [27] Weidmann J., *Linear Operators in Hilbert Spaces*, Springer-Verlag, Berlin, 1980.
- [28] Zhislin G.M., A study of the spectrum of the Schrödinger operator for a system of several particles, *Trudy Moskov. Mat. Obsc.* **9**, 81-120 (1960).

D.H. JAKUBASSA-AMUNDSEN, MATHEMATICS INSTITUTE, UNIVERSITY OF MUNICH, THERE-  
SIENSTR. 39, 80333 MUNICH, GERMANY  
E-mail address: [dj@mathematik.uni-muenchen.de](mailto:dj@mathematik.uni-muenchen.de)